

## Healthcare workers' radiation exposure tied to range of health problems

*Date:* April 12, 2016

*Source:* American Heart Association

*Summary:* Healthcare workers in cardiac catheterization labs may have higher odds than workers elsewhere for health problems including cataracts, skin lesions, cancers and orthopedic illness. Estimated exposure to radiation was highest for interventional cardiologists and electrophysiologists. Intensive training in radiation protection can reduce occupational doses dramatically.

### FULL STORY

---

Healthcare professionals performing x-ray guided cardiovascular procedures may be at higher risk for health problems including orthopedic problems, cataracts, skin lesions and cancers, according to new research in the American Heart Association journal *Circulation: Cardiovascular Interventions*.

The study identified potential radiation-related health risks to professionals that perform common cardiovascular procedures using fluoroscopy -- an imaging technique that uses X-rays to obtain real-time moving images of the heart. Such procedures include coronary angiography to diagnose heart problems and coronary artery angioplasty to widen narrowed arteries. These procedures are commonly done in a hospital cardiac catheterization (cath) lab.

"For experienced, busy interventional cardiologists and electrophysiologists, annual exposure to radiation is around 5 millisievert (mSv, a unit of measure for radiation's biological effects)," said Maria Grazia Andreassi, M.Sc., Ph.D., lead author of the study and head of the Genetics and Molecular Epidemiology Unit at the National Research Council Institute of Clinical Physiology in Pisa, Italy. That amount adds up to 50 to 200 mSv, or the equivalent of 2,500 to 10,000 chest X-rays, over a 30-year career.

Compared to healthcare professionals not exposed to radiation, cath lab workers (including doctors, nurses and technicians) who were exposed to radiation for a median of 10 years had:

2.8 times higher odds of having skin lesion; 7.1 times higher odds of having orthopedic (back/neck/knee) problems; and 6.3 times higher odds of having cataracts.

In addition, among cath lab workers, estimated radiation exposure was highest for the interventional cardiologists and electrophysiologists. The odds of having medical problems also increased for workers with more than 16 years of history in the cath lab. After adjusting for age, gender and smoking, workers with more than 16 years of history in the cath lab also had 3 times high odds of having cancer.

Cath lab workers also had elevated rates of high blood pressure and high cholesterol, the study found, but relatively low rates of cardiovascular illness. However, previous epidemiological studies have linked chronic radiation exposure with long-term cardiovascular effects, Andreassi said.

"Occupational doses of radiation in cardiovascular procedures guided by fluoroscopy are the highest doses registered among medical staff using X-rays," Andreassi said. "Interventional cardiologists and electrophysiologists have a two to three times higher annual exposure than that of radiologists, as they are closer to the radiological source and experience radiation exposure with the patient, whereas diagnostic radiologists are generally shielded from radiation exposure."

The new findings are based on an Italian survey answered by 466 workers who had an average of 10 years' experience working in cardiac cath labs, plus 280 professionals who worked in other settings. Findings were adjusted for the fact that the cath lab workers were more likely to be smokers, as well as other relevant factors, Andreassi said.

Noting that intensive training in radiation protection can reduce occupational doses dramatically, Andreassi said every effort should be made to raise radiation awareness in the professional communities of interventional cardiologists and electrophysiologists. "Unfortunately, cardiologists pay little heed to monthly or cumulative reports of radiation exposure. And recent studies confirm that simple, effective protection measures -- such as a lead curtain, protection glasses and thyroid collars -- are not used by the majority of exposed cardiologists."

Study limitations noted by researchers included: survey respondents themselves chose whether to participate (and if they had health problems, may have been more motivated to do so) and radiation doses were self-reported, not directly measured. A direct cause-and-effect link between radiation exposure and health problems cannot be inferred from the results of this study.

---

### Story Source:

The above post is reprinted from materials provided by **American Heart Association**. *Note: Materials may be edited for content and length.*

---

### Journal Reference:

1. Maria Grazia Andreassi, Emanuela Piccaluga, Giulio Guagliumi, Maurizio Del Greco, Fiorenzo Gaita, and Eugenio Picano. **Occupational Health Risks in Cardiac Catheterization Laboratory Workers**. *Circulation: Cardiovascular Interventions*, April 2016 DOI: 10.1161/CIRCINTERVENTIONS.115.003273

---

### Cite This Page:

MLA

APA

Chicago

---

American Heart Association. "Healthcare workers' radiation exposure tied to range of health problems." ScienceDaily. ScienceDaily, 12 April 2016.  
<[www.sciencedaily.com/releases/2016/04/160412211121.htm](http://www.sciencedaily.com/releases/2016/04/160412211121.htm)>.

---

## Recommended Articles

Physicians' exposure to radiation prompt cellular changes that may protect the body from harm

European Society of Cardiology, ScienceDaily, 2011

Real-time radiation monitor can reduce radiation exposure for medical workers

UT Southwestern Medical Center, ScienceDaily, 2014

Cardiologists dramatically cut patient radiation exposure from X-rays

Mayo Clinic, ScienceDaily, 2012

Improved imaging takes X-ray risks out of the picture

University of Missouri-Columbia, ScienceDaily, 2016

Working in an interventional laboratory may lead to health problems

American College of Cardiology, ScienceDaily, 2015

Radiation exposure increases risk of health problems for medical staff  
Honor Whiteman, Medical News Today

Cumulative Radiation Exposure From Imaging Scans Should Be Weighed Against The Benefits Say Researchers  
Catharine Paddock PhD, Medical News Today, 2009

Pause and pulse: radiation dose in pediatric fluoroscopy.

Beverley Newman et al., *Pediatr Rev*, 2011

Cumulative effective doses from radiologic procedures for pediatric oncology patients.

Bilal A Ahmed et al., *Pediatrics*, 2010

Radiation Exposure and Cancer Risk in Congenital Heart Disease

American Academy of Pediatrics, AAP Grand Rounds, 2014